



**DISTRIBUTION AND ABUNDANCE OF PEARY CARIBOU (*Rangifer tarandus pearyi*)  
ON LOUGHEED ISLAND, JULY 2016**

MORGAN ANDERSON<sup>1</sup>

Version: 13 August 2016

<sup>1</sup>Wildlife Biologist High Arctic, Department of Environment  
Wildlife Research Section, Government of Nunavut Box 209 Igloolik NU X0A 0L0

STATUS REPORT 2016-02  
NUNAVUT DEPARTMENT OF ENVIRONMENT  
WILDLIFE RESEARCH SECTION  
IGLOOLIK, NU





## Contents

List of Figures.....	v
List of Tables.....	v
Introduction.....	6
ላይኛው ልማት ስራ.....	6
Study Area.....	7
Methods.....	7
<i>Aerial Survey</i> .....	7
<i>Analysis</i> .....	7
Results .....	9
የሥራው ውጤት.....	9
Discussion .....	11
የሥራው ጠቃሚነት.....	13
Management Recommendations .....	14
የሥራው ጠቃሚነት ለማረጋገጥ.....	14
Acknowledgements .....	14
Literature Cited.....	15
Appendix 1. Alternate population calculations. ....	16
<i>Jolly Method II Calculations</i> .....	16







population calculations following Jolly's Method II are provided in Appendix 4, along with calculations following a systematic stratified survey design (Cochran 1977). The muskoxen and caribou detected in this survey were patchily distributed and serially correlated, not randomly distributed. For systematic samples from serially correlated populations, estimates of uncertainty based on deviations from the sample mean are expected to be upwardly biased and influenced by the degree of serial correlation; high serial correlation implies that there is less random variation in the unsurveyed sections between systematically spaced transects than if serial correlation were low (Cochran 1977). Calculating uncertainty based on nearest-neighbor differences incorporates serial correlation, and the upward bias in the uncertainty is expected to be less than if it were calculated based on deviations from the sample mean. Nearest-neighbor methods have been used previously to calculate variance around survey estimates on the unweighted ratio estimate (Kingsley et al. 1981, Stirling et al. 1982, Kingsley et al. 1985, Anderson and Kingsley 2015).

The model for observations on a transect survey following Cochran (1977) is:

$$y_i = Rz_i + \varepsilon_i \sqrt{z_i}$$

Where  $y_i$  is the number of observations on transect  $i$  of area  $z_i$ ,  $R$  is the mean density and error terms  $\varepsilon_i$  are independently and identically distributed. In this model, the variance of the error term is proportional to the area surveyed. The best estimate of the mean density  $\hat{R}$  is:

$$\hat{R} = \frac{\sum_i y_i}{\sum_i z_i}$$

The error sum of squares, based on deviations from the sample mean, is given by:

$$\left( \sum_i \frac{y_i^2}{z_i} \right) - \frac{(\sum_i y_i)^2}{\sum_i z_i}$$

The finite-population corrected error variance of  $\hat{R}$  is:

$$Var(\hat{R}) = \frac{(1-f)}{(n-1)\sum_i z_i} \left( \left( \sum_i \frac{y_i^2}{z_i} \right) - \frac{(\sum_i y_i)^2}{\sum_i z_i} \right)$$

Where  $f$  is the sampling fraction and  $n$  is the number of transects. The sampling fraction also provides the scaling factor for moving from a ratio (population density) to a population estimate. It is calculated as  $(\sum z_i)/Z$ , where  $Z$  is the study area and  $\sum z_i$  is the area surveyed. The irregular study area boundaries mean that  $f$  varies from the 20% sampling fraction expected from a 1-km survey strip and 5-km transect spacing.

If we were to apply a model  $y_i = Rz_i + \varepsilon_i$  instead, then the variance of the error term would be independent of  $z$ , so the variance would depend on the number of items in the sample, but not their total size. This would lead to a least squares estimate of  $R$  of  $\sum zy / \sum z^2$ , rather than the more intuitive density definition and model for  $R$  presented above.

To incorporate serial correlation in the variance, we used a nearest-neighbor calculation, with the error sum of squares given by:









Island, at least 110 km across the sea ice from Bathurst Island (Poole et al. 2015). She then continued 110 km across the ice to Borden Island, where she died in December 1995 (Poole et al. 2015).




Figure 3. Population estimates for Peary caribou on Loughheed Island. Grey bars indicate estimates including calves (Tener 1963, Miller et al. 1977, this report), black bars are minimum counts (Miller et al. 1977, Miller 1987, this report for 2015), and white bars are population estimates of 1+-year-old caribou (Gunn and Dragon 2002, Jenkins et al. 2011).

Although not conducted as a survey, we did fly over Loughheed Island in 2015 to determine whether we could collect pellet samples using a Twin Otter drop-off and pick-up, or whether a helicopter would be required. We counted at least 119 Peary caribou during the flight, including some groups of 15-20 individuals (in which case the lower value was added for the minimum count of 119; Figure 4). Flight height was 90-150 m above ground and conditions were clear and sunny, with one observer each side of the plane and a navigator/recorder. No marks were made on the wing struts to define a survey strip.





Shelf Program for logistical support. Funding for this work was provided by the Nunavut Department of Environment Wildlife Research Section. The Polar Continental Shelf Program provided logistical support (Project 308-16). The work was completed under Government of Nunavut Wildlife Research Section permit 2016-010.

### Literature Cited

- Anderson, M. and M. C. S. Kingsley. 2015. Distribution and abundance of Peary caribou (*Rangifer tarandus pearyi*) and muskoxen (*Ovibos moschatus*) on southern Ellesmere Island, March 2015. Nunavut Department of Environment, Wildlife Research Section, Status Report, Igloodik, NU. 46 pp.
- Caughley, G. 1977. Sampling in aerial survey. *Journal of Wildlife Management* 41(4): 605-615.
- Cochran, W. G. 1977. Sampling techniques. 3<sup>rd</sup> ed. Wiley, New York, NY. 428 pp.
- Gould, W. A., M. Reynolds, and D. A. Walker. 2003. Vegetation, plant biomass, and net primary productivity patterns in the Canadian Arctic. *Journal of Geophysical Research* 108 (D2): 8167, doi:10.1029/2001JD000948.
- Jenkins, D., M. Campbell, G. Hope, J. Goorts, and P. McLoughlin. 2011. Recent trends in abundance of Peary Caribou (*Rangifer tarandus pearyi*) and muskoxen (*Ovibos moschatus*) in the Canadian Arctic Archipelago, Nunavut. Department of Environment, Government of Nunavut, Wildlife Report No. 1, Pond Inlet, Nunavut. 184 pp.
- Jolly, G. M. 1969. Sampling methods for aerial censuses of wildlife populations. *East African Agricultural and Forestry Journal* 34 (special issue):46-49.
- Kingsley, M. C. S., and G. E. J. Smith. 1981. Analysis of data arising from systematic transect surveys. Pp 40-48 in F. L. Miller and A. Gunn, eds. Proceedings, Symposium on Census and Inventory Methods for Populations and Habitats, Banff, AB, April 1980.
- Kingsley, M. C. S., I. Stirling, and W. Calvert. 1985. The distribution and abundance of seals in the Canadian high arctic, 1980-82. *Canadian Journal of Fisheries and Aquatic Sciences* 42 (6): 1189-1210.
- Miller, F. L., R. H. Russell, and A. Gunn. 1977. Distributions, movements and numbers of Peary caribou and muskoxen on western Queen Elizabeth Islands, Northwest Territories, 1972-74. *Canadian Wildlife Service Report Series* No. 40, Edmonton, Alberta. 55 pp.
- Miller, F. L. 1987. Peary caribou and muskoxen on Bathurst, Alexander, Marc, Massey, Vanier, Cameron, Helena, Loughheed, and Edmund Walker islands, Northwest Territories, July 1985. Technical Report Series No. 20, Canadian Wildlife Service, Western and Northern Region, Alberta. 45 pp.
- Miller, F. L. and S. J. Barry. 2009. Long-term control of Peary caribou numbers by unpredictable, exceptionally severe snow or ice conditions in a non-equilibrium grazing system. *Arctic* 62(2): 175-189.
- Miller, F.L., and A. Gunn. 2003. Catastrophic die-off of Peary caribou on the western Queen Elizabeth Islands Canadian High Arctic. *Arctic* 56: 381–390.
- Miller, F. L., R. H. Russell and A. Gunn. 1975. The decline of caribou on the western Queen Elizabeth Islands. *Polarforschung* 45: 17-22.
- Norton-Griffiths, M. 1978. Counting animals. In: J. J. R. Grimsdell, ed. *Handbooks on techniques currently used in African wildlife ecology*. No 1. AWLF, Nairobi, Kenya.
- Poole, K. G., A. Gunn, J. Wierzchowski, and M. Anderson. 2015. Peary caribou distribution within the Bathurst Island Complex relative to the boundary proposed for Qausuittuq National Park, Nunavut. *Rangifer* 35 Special Issue 23: 81-98.
- Stirling, I., M. C. S. Kingsley and W. Calvert. 1982. The distribution and abundance of seals in the Beaufort Sea, 1974-1979. *Canadian Wildlife Service Occasional Paper* 47.
- Tener, J. S. 1963. Queen Elizabeth Islands game survey, 1961. *Canadian Wildlife Service Occasional Paper* No. 4. 50 pp.
- Walker, D. A., M. K. Reynolds, F. J. A. Daniëls, E. Einarsson, A. Elvebakk, W. A. Gould, A. E. Katenin, S. S. Kholod, C. J. Markon, E. S. Melnikov, N. G. Moskalenko, S. S. Talbot, B. A. Yurtsev, and the other members of the CAVM Team. 2005. The Circumpolar Arctic vegetation map. *Journal of Vegetation Science* 16: 267-282.

**Appendix 1. Alternate population calculations.**

*Jolly Method II Calculations*

In this report, we used a systematic sampling approach to analysis, since we were estimating abundance of a patchy population rather than estimating density in a habitat (which varied across the study area). Other systematic aerial surveys have frequently used Jolly’s Method II, and estimates derived from both analyses were similar. Population estimates for fixed-width strip sampling using Jolly’s Method 2 for uneven sample sizes (Jolly 1969; summarized in Caughley 1977) are derived as follows:

$$\hat{Y} = RZ = Z \frac{\sum_i y_i}{\sum_i z_i}$$

Where  $\hat{Y}$  is the estimated number of animals in the population,  $R$  is the observed density of animals (sum of animals seen on all transects  $\sum_i y_i$  divided by the total area surveyed  $\sum_i z_i$ ), and  $Z$  is the total study area. The variance is given by:

$$Var(\hat{Y}) = \frac{N(N - n)}{n} (s_y^2 - 2Rs_{zy} + R^2s_z^2)$$

Where  $N$  is the total number of transects required to completely cover study area  $Z$ , and  $n$  is the number of transects sampled in the survey.  $s_y^2$  is the variance in counts,  $s_z^2$  is the variance in areas surveyed on transects, and  $s_{zy}$  is the covariance. The estimate  $\hat{Y}$  and variance  $Var(\hat{Y})$  are calculated for each stratum and summed. The Coefficient of Variation ( $CV = \sigma/\hat{Y}$ ) was calculated as a measure of precision.

Table 4. Abundance estimates (Jolly 1969 Method II) for caribou on Lougheed Island, July 2016.  $N$  is the total number of transects required to completely cover study area  $Z$ ,  $n$  is the number of transects sampled in the survey covering area  $z$ ,  $y$  is the observed muskoxen,  $Y$  is the estimated muskoxen with variance  $Var(Y)$ . The coefficient of variation (CV) is also included.

<b>Y</b>	<b>Var(Y)</b>	<b>n</b>	<b>Z (km<sup>2</sup>)</b>	<b>z (km<sup>2</sup>)</b>	<b>N</b>	<b>y</b>	<b>Density (per km<sup>2</sup>)</b>	<b>CV</b>
140	1511.91	4	1359.58	252.13	24	26	0.1031	0.28